



LIQUID BIOCELL[®] LIFE

A Scientific Perspective

Liquid BioCell[®] Life is a delicious liquid nutraceutical that promotes healthy aging, active joints, and younger-looking skin. It provides a unique and proprietary combination of science-substantiated ingredients and antioxidant-rich phytonutrients that synergistically support a healthier life. Only Liquid BioCell[®] Life contains multi-patented Liquid BioCell[®], containing constituents such as collagen, hyaluronic acid (HA), and chondroitin sulfate (CS), which are essential for healthy joints and skin. It is blended with 13 exotic, antioxidant-rich fruits from around the globe, including blueberry, mangosteen, açai berry, noni, and pomegranate. The phytonutrients contained in the fruits possess multiple health-promoting, biological activities.

Maintaining active joints and younger-looking skin is essential for healthy aging. The key ingredient is Liquid BioCell[®], a patented dietary supplement proven to improve joint and skin health. Its constituents, type II collagen (hydrolyzed), hyaluronic acid, and chondroitin sulfate (depolymerized) are proteins and glycosaminoglycans (GAGs) critical for the health of these tissues. Other key ingredients are resveratrol and 13 superfruits carefully selected from around the world to provide a variety of phytonutrients, including vitamins, flavonoids, polyphenols, and dietary fibers. The flavor is so unique, you will remember exactly where you were the first time Liquid BioCell[®] Life touched your palate.

Studies

Liquid BioCell[®] Life contains a number of health-promoting, functional ingredients. They include patented Liquid BioCell[®], resveratrol, and 13 superfruits known to contain antioxidants, vitamins, minerals and dietary fibers. Scientific studies on individual ingredients suggest that it can synergistically promote the overall health of aging people.


Safety

Several toxicology studies, including a 14-day acute study and a 90-day sub-chronic study, have demonstrated that Liquid BioCell[®] is well tolerated up to over 30 times the recommended human dose of 30 mg/kg body weight/day (1). In a pilot clinical trial involving humans with joint health conditions, Liquid BioCell[®] (2g/day) was also well tolerated during the 8-week study period (2). Resveratrol is not known to cause adverse effects in humans, as a recent trial that evaluated the safety of oral resveratrol in 10 healthy volunteers found a single dose up to 5 grams resulted in no serious adverse effects (3).

Joint Health

Studies support that the functional nutrients in Liquid BioCell[®] Life can help improve joint health by providing structurally and functionally essential molecules and also by protecting the joints from oxidative damage.


There are many different types of collagen found in the body, and type II is the most abundant (>90% of collagens) in the cartilage of the synovial joint, the most common and movable joint (4). Type II collagen forms a fibrillar network to provide tensile strength to the cartilage. Proteoglycans (PGs) such as aggrecans are another key component of articular cartilage, and comprise a core protein linked with various GAGs such as HA and CS (5).



◀ This diagram demonstrates how Liquid BioCell® mirrors the joint composition, a key reason why the body readily accepts it.


PGs are immobilized into the collagen fibrillar network to provide resistance to compression. The contents of these structural components and the integrity of the cartilage matrix decrease due to aging or degenerative conditions. Liquid BioCell®'s composition in terms of type II collagen and proteoglycans mirrors that of human articular cartilage and can support healthy joints by supplementing these molecules (Figure 1).

Figure 1. Comparison of composition of Liquid BioCell™ and human articular cartilage.



Multiple clinical studies demonstrate the safety and beneficial effects of collagen and GAGs toward various joint conditions (6; 7; 8). A double-blind, placebo-controlled trial of Liquid BioCell® also revealed statistically significant efficiency in supporting joint health, leading to a decrease in difficulty carrying out physical activities (Figure 2) (2).


Figure 2. Reduced difficulty in physical activities 8-weeks after daily ingestion of Liquid BioCell®.



Recent studies suggest several possible mechanisms of action for the joint health benefits provided by Liquid BioCell®. First, it may support the cartilage matrix of collagen fibrillar network filled with proteoglycans by providing constituent molecules, and also supplement joint fluid by providing HA, which helps lubricate the joint (9). A review published in Current Medical Research and Opinion comprehensively analyzed collagen hydrolysate studies listed in the PubMed database (8). The authors stated, “According to published research, orally administered collagen hydrolysate has been shown to be absorbed intestinally and to accumulate in cartilage.”


In order to ensure fast absorption and higher bioavailability, the three key constituents of Liquid BioCell® are properly processed. Type II collagen is hydrolyzed to have an average molecular weight of 1.3 to 1.8 kDa, while chondroitin sulfate is depolymerized. In an HA bioavailability study, subjects taking Liquid BioCell® (1500 mg/daily) for 28 days were shown to increase steady-state HA levels in their blood as much as 60-fold (Figure 3).

Figure 3. Increase in steady-state HA levels.



Second, a recent finding of the ability of hydrolyzed collagen, but not undenatured collagen, to stimulate novel biosynthesis of type II collagen from chondrocytes suggests that hydrolyzed collagen in Liquid BioCell® may directly stimulate the only resident cells in the cartilage to produce its matrix components (Figure 4) (10). This would help maintain the tensile strength of cartilage, thereby supporting the shock absorption capability of the joint.


4. Figure 4. Hydrolyzed collagen in Liquid BioCell® may directly stimulate the only resident cells in the cartilage to produce its matrix components.



Joint support by Liquid BioCell® is augmented by various fruit-derived phytonutrients with potent antioxidant activity. These phytochemicals contained in Liquid BioCell® Life have been shown to have various health-promoting biological properties including antioxidant activity. Research suggests that antioxidants can support normal functions of chondrocytes, which appear to be affected by oxidative stress in the joints. First, nitrotyrosine, a marker of oxidative stress, is over-expressed in aging or degrading cartilage (11). This oxidative stress can lead to growth arrest and senescence of chondrocytes (12;13), a manifestation of the loss of their normal function, causing in turn deterioration in the structure and function of the cartilage (14). Second, lower antioxidative capacity was observed in degraded regions as compared with the intact regions from the same cartilage. Furthermore, treatment of chondrocytes with reactive oxygen species (ROS) induced genome instability and reduction of replicative capacity and GAG production, whereas treatment with antioxidants such as ascorbic acid (vitamin C) prevented the damaging effects (15).

These studies suggest that high contents of antioxidants such as vitamins, flavonoids, and polyphenols in Liquid BioCell® Life may help neutralize these damaging effects of oxidative stress in aging joints. Taken together, a variety of naturally-occurring antioxidants abundantly contained in Liquid BioCell® Life can support joint health by neutralizing oxidative stress, in synergy with Liquid BioCell® in supporting normal functions of chondrocytes and cartilage matrix they produce.

5. Figure 5. Dose-dependent inhibition of hyaluronidase



Skin health

The skin, the largest organ in the body, is the epidermis and the dermis. The dermis shares key structural components with the joint, including hyaluronic acid (HA), chondroitin sulfate (CS) and collagen (types I and III vs type II in the joint). Thus, Liquid BioCell® Life helps support healthy and younger-looking skin through the provision of structurally-essential molecules to the dermis and through protection against oxidative damage closely associated with intrinsic and extrinsic skin aging.

HA, a key component of Liquid BioCell®, is found in virtually all tissues including the skin and bodily fluids (16). Approximately one-third of HA is turned over daily by coordinated action of HA synthesis and hyaluronidases (17). HA in the skin is a major component of the dermal extracellular matrix and responsible for hydration, nutrient exchange, protection against free radical damage, and regulation of cell differentiation and motility (18). As we grow older, the contents of dermal structural components decrease, resulting in disorganization or collapse of the extracellular network in the dermis (19). The molecular weight of HA also decreases, aggravating the loss of its water-retaining capability.

Studies suggest that Liquid BioCell® can help maintain volume, density, and turgidity of the skin in several ways. First, Liquid BioCell® harbors a unique biological activity which differentiates it from other dietary ingredients and products for skin health.

Second, a bioavailability study (Figure 3) suggests that a 60-fold increase in blood HA levels from daily intake of Liquid BioCell® provides a pool available for the maintenance of adequate HA levels in the skin. Photoaging processes are similar to those of intrinsic, chronological aging. Dai et al (2007) showed that during photoaging, the total cell number and percentage of proliferating fibroblasts in the papillary dermis decreased in concomitance with the loss of HA from the dermis (20). It may alleviate the effects of photoaging by supplementing HA.

Third, the HA of Liquid BioCell® may help skin health in a more fundamental way. Greco et al reported that during wound healing, HA stimulated the proliferation of dermal fibroblasts responsible for biosynthesis of collagen, elastin and GAGs including HA (21; 22). Enhanced bioavailability of HA and inhibition of hyaluronidase would increase HA levels in the skin, which is postulated to stimulate dermal fibroblasts to synthesize skin collagens and other extracellular components.

Hyaluronic acid and collagen are naturally-occurring elements in human skin tissue and are vital for maintaining youthful skin. Used safely and effectively worldwide for years, HA can be described as a “space filler” because it provides the moisture-filled cushioning in between our cells, while collagen provides the connective framework to uphold the skin tissue’s integrity. HA is especially abundant in youthful skin, where it binds to water and provides volume — thereby plumping out the appearance of wrinkles and adding resiliency to skin.

Healthy aging requires an effective measure against oxidative stress in the skin because insufficient removal of reactive oxygen species (ROS) initiates or accelerates both natural and extrinsic skin aging processes by damaging DNA, proteins and cellular membrane (23;12). Our body has endogenous defense mechanisms, including enzymatic (superoxide dismutase and catalase, etc.) and nonenzymatic molecules (vitamins and glutathione) (24). As one ages, however, these defense mechanisms weaken while the load of oxidative stress increases, which will accelerate skin aging. Thus, additional and constant supply of antioxidants is essential.

As shown in Table 1, 13 fruits contained in Liquid BioCell® Life provide a variety of antioxidants, such as vitamins and polyphenols that can help neutralize ROS arising from normal cellular metabolism and from exogenous factors, such as UV exposure, cigarette smoking, and alcohol consumption (25; 26).



Table 1.

Fruits	Phytochemical Compounds
Blueberry	Anthocyanins, flavonols, proanthocyanidins, resveratrol
Red Grapes	Anthocyanins, resveratrol, flavonoids
Strawberry	Anthocyanins, ellagic acid
Apple	Quercetin, chlorogenic acid, epicatechin, procyanidin B2
Pomegranate	Punicalagins, catechins, anthocyanins, quercetin
Mangosteen	Mangostins
Acai Berry	Carotenoids, anthocyanins, flavonols
Jujube	Triterpenes, cyclopeptide alkaloids, flavonoids
Lyium (Ozai berry)	LBP (Lycium Barbarum polysaccharide)
Nopal	Prickly Pear Pectin
Maqui	Anthocyanins, flavonols
Noni	Lignans, flavonoids, catechins
Cranberry	Proanthocyanidins

Phytonutrients in Liquid BioCell® Life possess a wide range of biological activities

Studies show that strawberries and oranges are rich in vitamin C, a powerful antioxidant that can only be obtained from food. Sunlight and environmental pollution can deplete vitamin C in the skin, causing lipid peroxidation of epidermal layers (26). Vitamin C plays another important role in maintaining healthy skin, as a co-factor required for collagen biosynthesis from dermal fibroblasts (27). Liquid BioCell® Life offers a continuous supply of vitamin C and various antioxidants, together with structural molecules such as HA essential to skin health, all of which can help neutralize the detrimental effects of sun exposure and associated aging process.

The antioxidant capacity of Liquid BioCell® Life is fortified further by carefully selected fruits from around the globe that harbor science-substantiated phytonutrients. For example, punicalagins and its hydrolysis product ellagic acid, the most abundant polyphenolic compound in pomegranate, have been shown to have potent antioxidant capacity (28). Studies also show that pomegranate extract restores enzymatic antioxidant defense mechanisms, including catalase, peroxidase and superoxide dismutase, which might explain pomegranate-induced amelioration of UVA-mediated damages and protection against the adverse effects of UVB radiation in human epidermal keratinocytes (29; 30). Similarly, anthocyanin-rich blueberry extract was shown to dampen UVB-mediated toxicity in human dermal fibroblasts through blocking NF- κ B and p53 activation and through suppressing ROS-mediated expression of matrix metalloproteinases (MMPs) that could degrade skin collagen (31). These results support that polyphenolic compound-rich pomegranate, blueberry and other fruits in Liquid BioCell® Life may be highly protective against UV-induced skin photoaging, as well as chronological aging.

Moreover, resveratrol, which is implicated in the “French Paradox,” is also found in Liquid BioCell® Life. Each serving contains the same amount of resveratrol as four full bottles of red wine. Studies showed that it had diverse biological properties in-vitro and prevented UVB damage and abnormal cell proliferation in skin whose effects appeared to be mediated through modulation of surviving (32; 33). Resveratrol administration increased the lifespans of yeast, worms, fruit flies, fish, and mice fed a high-calorie diet (34), and intensive research is underway to investigate whether resveratrol has similar longevity effects in humans. Interestingly, in a recent aging study, a low dose of dietary resveratrol had altered gene expression in the heart, brain, and skeletal muscle similar to that induced by caloric restriction (35).

REFERENCES SUPPORTING LIQUID BIOCELL® LIFE and its constituents

1. Acute and subchronic oral toxicity studies in rats of a hydrolyzed chicken sternal cartilage preparation. Schauss et al. 2007, *Food Chem Toxicol*, pp. 45: 315-321.
2. A randomized double-blind clinical pilot trial evaluating the safety and efficacy of hydrolyzed collagen type II (BioCell Collagen® II) in adults with osteoarthritis. Sheldon et al. 2003, Miami Research Associates.
3. Boocock et al. Phase I dose escalation pharmacokinetic study in healthy volunteers of resveratrol, a potential chemopreventive agent. *Cancer Epidemiol Biomarkers Prev*. 2007, Vols. 16(6): 1246-52.
4. Articular cartilage: structure, injuries and review of management. Bhosale et al. 2008, *Brit Med Bull*, pp. 87: 77-95.
5. Aggrecan, aging, and assembly in articular cartilage. Dudhia. 2005, *Cell MolLife Sci*, pp. 62: 2241-2256.
6. Effects of Oral Administration of Type II Collagen on Rheumatoid Arthritis. Trentham et al. 1993, *Science*, pp. 261: 1727-1730.
7. Suppression of type II collagen-induced arthritis by the intravenous administration of type II collagen or its constituent peptide. Englert et al. 1984, *Cell Immuno*, pp. 87: 357-365.
8. Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: a review of the literature. Bello et al. 2006, *Curr Med Res Opin*, pp. 22(11): 2221-2232.
9. Hyaluronan synthesis and degradation in cartilage and bone. Bastow. 2008, *Cell Mol Life Sci*, pp. 65: 395-413.
10. Stimulation of type II collagen biosynthesis and secretion in bovine chondrocytes cultured with degraded collagen. Oesser et al. 2003, *Cell Tissue Res*, pp. 399: 393-399.
11. Detection of nitrotyrosine in aging and osteoarthritic cartilage: Correlation of oxidative damage with the presence of interleukin-1beta and with chondrocyte resistance to insulin-like growth factor 1. Loeser et al. 2002, *Arthritis Rheum*, pp. 46: 2349-2357.
12. Oxidants, oxidative stress, and the biology of aging. Finkel et al. 2000, *Nature*, pp. 408: 239-247.
13. Potential involvement of oxidative stress in cartilage senescence and development of osteoarthritis: oxidative stress induces chondrocyte telomere instability and down-regulation of chondrocyte function. Yudoh et al. 2005, *Arthritis Res Ther*, pp. 7: R380-R391.
14. Telomere erosion and senescence in human articular cartilage chondrocytes. Martin et al. 2001, *J Gerontol A Biol Sci Med Sci*, pp. 56: B172-B179.
15. Loeser. Aging and osteoarthritis: the role of chondrocyte senescence and aging changes in the cartilage matrix. *Osteoarthritis and Cartilage*. 2009, Vols. 17:971-979.
16. Hyaluronic acid in cutaneous intrinsic aging. Ghersetich et al. 1994, *Int J Dermatol*, pp. 33(2): 119-122.
17. Stern et al. Hyaluronan catabolism: a new metabolic pathway. *Eur J Cell Biol*. 2004, Vols. 83(7): 317-325.
18. Wiest et al. Native hyaluronic acid in dermatology-results of an expert meeting. *J Dtsch Dermatol Ges*. 2008, Vols. 6(3): 176-180.
19. Collagen fragmentation promotes oxidative stress and elevates matrix metalloproteinase-1 in fibroblasts in aged human skin. Fisher et al. 2009, *Am J Pathol*, pp. 174: 101-114.
20. Chronic ultraviolet B irradiation causes loss of hyaluronic acid from mouse dermis because of down-regulation of hyaluronic acid synthesis. Dai et al. 2007, *Am J Pathol*, pp. 282: 1451-1561.
21. Hyaluronic acid stimulates human fibroblast proliferation within a collagen matrix. Greco et al. 1998, *J Cell Physiol*, pp. 177(3): 465-473.
22. Hyaluronan facilitates transforming growth factor-beta1-mediated fibroblast proliferation. Meran et al. 2008, *J Biol Chem*, pp. 283: 6530-6545.
23. Aging: a theory based on free radical and radiation chemistry. Harman. 1956, *J Gerontol*, pp. 11(3): 298-300.
24. Enzymatic and nonenzymatic antioxidants in epidermis and dermis of human skin. Shindo et al. 1994, *J Invest Dermatol*, pp. 102(1): 122-124.
25. Intake of flavonoids and flavones and risk of coronary heart disease in male smokers. Hirvonen et al. 2001, *Epidemiology*, pp. 12: 62-67.
26. In vivo exposure to ozone depletes vitamins C and E and induces lipid peroxidation in epidermal layers of murine skin. Thiele et al. 1997, *Free Radic Biol Med*, pp. 23(3): 385-91.
27. Ascorbic acid specifically increases type I and type III procollagen messenger RNA levels in human skin fibroblast. Geesin et al. 1988, *J Invest Dermatol*, pp. 90(4): 420-424.
28. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. Gil et al. 2000, *J Agric Food Chem*, pp. 48(10): 4581-4589.
29. Studies on antioxidant activity of pomegranate (*Punica granatum*) peel extract using in vivo models. Chidambara et al. 2000, *J Agric Food Chem*, pp. 50(17): 4791-4795.
30. Photochemo preventive effect of pomegranate fruit extract on UVA-mediated activation of cellular pathways in normal human epidermal keratinocytes. Syed et al. 2006, *Photochem Photobiol*, pp. 82(2): 398-405.

31. Bog blueberry anthocyanins alleviate photoaging in ultraviolet-B irradiation-induced human dermal fibroblasts. Bae et al. 2009, Mol Nutr Food Res, pp. 53(6): 726-738.
32. Prevention of ultraviolet-B radiation damage by resveratrol in mouse skin is mediated via modulation in survivin. Aziz et al. 2005, Phytochem Photobiol, pp. 81(1): 25-31.
33. Chemoprevention of skin cancer by grape constituent resveratrol: relevance to human disease? Aziz et al. 005, FASEB J, pp. 19(9): 1193-1195.
34. A multitargeted agent for age-associated chronic diseases. Kuzhuvellil et al. 2008, Cell Cycle, p. 7:8.
35. A low dose of dietary resveratrol partially mimics caloric restriction and retards aging parameters in mice. Barger et al. 2008, PLoS One, p. 3(6): e2264.
36. Schauss et al. 2012. Effect of the novel low molecular weight hydrolyzed chicken sternal cartilage extract, Liquid BioCell, on improving osteoarthritis-related symptoms: a randomized, double-blind, placebo-controlled trial. J Agric Food Chem. 60(16): 4096-101.
37. Schwartz et al. 2012. Ingestion of Liquid BioCell, a novel hydrolyzed chicken sternal cartilage extract, enhanced blood microcirculation and reduced facial aging signs. Clin Interv Aging. 7: 267-273.

* Liquid BioCell is the exclusive highly bioavailable liquid form of BioCell Collagen® TF

FDA Disclaimer

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Legal Disclaimer

The information contained in this document (including but not limited to literature, abstracts, clinical studies, research data, historical uses, advertisements, publications and any other information that is provided in the form of email or attachment) is solely intended for the purposes of education and knowledge, and should not be used as medical advice. Jusuru International, Inc. ("Jusuru") will not assume responsibility for how you use this information in the future. Not all of the clinical studies referenced and contained in this document are on Liquid BioCell® Life; the other studies are included as background information. Jusuru is not making any legal representation as to the information provided herein. While efforts have been made to ensure the accuracy of the content and information contained herein, Jusuru gives no warranty as to the accuracy of the information contained in the content of this document. Jusuru reserves the right to withdraw or delete information at any time.

Copyright Notice

All content included in this document, such as text, graphics, logos, images and data compilations is the sole property of Jusuru or its content suppliers and/or vendors and protected by United States and International Copyright Laws. The compilation of all content in this document is the exclusive property of Jusuru and protected by U.S. and International Copyright Laws.

Trademarks

Jusuru®, Liquid BioCell® Life (Logo and Design); and other marks indicated in our document are registered, or are currently under registration of trademarks of Jusuru International, Inc., or its subsidiaries, in the United States and other countries. Jusuru's graphics, logos, scripts, and service names are trademarks or trade dress of Jusuru or its subsidiaries. Jusuru's trademarks and trade dress may not be used in connection with any product or service that is not Jusuru in any manner that is likely to cause confusion among customers or in any manner that disparages or discredits Jusuru.

All other trademarks not owned by Jusuru or its subsidiaries that appear in this document are the property of their respective owners, who may or may not be affiliated with, connected to, or sponsored by Jusuru or its subsidiaries. Liquid BioCell® (Logo and Design) is a registered trademark of BioCell Technology, LLC.

Applicable Law

You agree that the laws of the State of California, without regard to principles of conflict of laws, will govern these Conditions of Use and any dispute of any sort that might arise between you and Jusuru or its affiliates.